



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF PENNSYLVANIA

TREE-äge™

EPA SLN No. PA-090001

**For control of Emerald Ash Borer in Ash Trees (*Fraxinus spp.*),
Tree Injection Only**

Active Ingredient:	
Emamectin Benzoate ¹	4.0%
Other Ingredients:	96.0%
Total:	100.0%

¹CAS No.155569-91-8

KEEP OUT OF REACH OF CHILDREN.

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use on label and in in booklet

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL
MAY RESULT IN POOR INSECT CONTROL, AND/OR CROP INJURY.**

PRECAUTIONARY STATEMENTS

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- long-sleeved shirt and long pants
- chemical resistant gloves (Category C) such as barrier laminate, butyl rubber >14 mils, nitrile rubber >14 mils, or neoprene rubber >14 mils.
- shoes and socks
- protective eyewear

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the TREE-äge container label.

This label must be in the possession of the user at the time of pesticide application.

APPLICATION TO ASH TREES (*Fraxinus* spp.)

TREE-äge is for control of emerald ash borer on ash trees growing in residential and commercial landscapes, parks, plantations, right of ways, seed orchards, and forested sites (in private, municipal, state, tribal and national areas). TREE-äge contains the active ingredient emamectin benzoate and is formulated to translocate in the tree's vascular system when injected. To assure optimum effectiveness, this product must be placed into active sapwood.

WHEN TO TREAT

TREE-äge contains the active ingredient emamectin benzoate which is a glycoside insecticide. It is active against larva and adult Emerald Ash Borer. The primary route of toxicity is through ingestion, but may also be lethal upon contact.

ENVIRONMENTAL CONDITIONS: Uptake of TREE-äge is dependent upon the tree's transpiration. Transpiration is dependent on a number of abiotic and biotic factors, such as soil moisture, soil and ambient temperature, and time of day. For optimal uptake, apply when soil is moist, soil temperatures are above 45°F, ambient temperatures are between 40° to 90°F, and during the 24 hour period when transpiration is greatest, typically before 2:00 PM. Applications to drought or heat stressed trees may result in injury to tree tissue, poor treatment and subsequent control. Watering the trees prior to injection may enhance the uptake of TREE-äge.

MONITOR TREE HEALTH and PEST INFESTATIONS: Effective injection treatment is favored by a full canopy (i.e., leaves) and healthy vascular system. Once these tissues are compromised by arthropod damage (larval galleries, defoliation, leaf mining, etc.) an effective and uniform application of TREE-äge may be difficult to achieve and subsequent control may be poor. Optimally, treatment should be made preventively at least 2 to 3 weeks before Emerald Ash Borer historically infest the host tree. As a result of systemic movement and longevity of TREE-äge in trees, this interval may be extended much earlier to 6 months should tree dormancy, adverse weather, management, asynchronous life cycle of pests, etc., allow earlier application timing.

TREE-äge may also be effective as a curative treatment against Emerald Ash Borer. Adult foliar feeding may be controlled within one month after treatment. During the larval stage, Emerald Ash Borer attacks the stem and branches and will disrupt vascular tissue that may result in poor distribution of TREE-äge in an infested tree. However, control may be achieved if larvae come into contact or feed on TREE-äge treated tissues.

USE

Use as formulated or dilute with equivalent 1 to 3 volumes of water or more, as necessary.

USE RATE TABLE

Tree Diameter (DBH) (Inches)	Low ml./tree	Medium ml./tree	Medium - High ml./tree	High ml./tree	Average No. Injection Sites*
4 to 6	15	25	50	-	3
7 to 9	20	40	80	-	4
10 to 12	30	55	110	165	5
13 to 15	35	70	140	210	6
16 to 18	40	75	150	225	7
19 to 21	50	100	200	300	8
22 to 24	-	115	230	345	10
25 to 27	-	130	260	390	11
28 to 30	-	145	290	435	12
31 to 33	-	160	320	480	13
34 to 36	-	175	350	525	15
37 to 39	-	190	380	570	16
40 to 42	-	205	410	615	17
43 to 45	-	220	440	660	18
46 to 48	-	235	470	705	20
49 to 51	-	250	500	750	21
52 to 54	-	265	530	795	22
55 to 57	-	280	560	840	23
58 to 60	-	295	590	885	25
61 to 63	-	310	620	930	26
64 to 66	-	325	650	975	27
67 to 69	-	340	680	1020	28
70 to 72	-	355	710	1065	30

* The number of injection sites listed is a guide for approximately how many are needed per size of tree.

For optimal control, it is recommended to be with ± 1 injection site of this number per tree. Higher rates tend to provide longer residual and control of more difficult to control insects. See **Target Pest** for additional information in choosing the amount of product to apply.

Applications in Trees

Tree Tissue to Protect	Target Pest	Recommended Rate	Comments
Foliage, Shoot, Stem, Trunk and Branch	Emerald Ash Borer- Adult and Larval stages	Low to High	For optimal control apply at least 30 days before historical egg hatch or adult flight and to trees whose vascular tissue is not damaged. If vascular tissue is damaged or plugged by insect galleries, nematodes or fungi, uniform treatment and control may not be achieved.

Compatibility

Do not mix TREE-äge before injection with other products such as insecticides, fungicides, plant growth regulators, surfactants, adjuvants, and fertilizers.

RESTRICTIONS

- Do not apply to trees that may yield food consumed by humans or used in animal feed.
 - Avoid treating trees that are moisture stressed or suffering from herbicide damage.
-

TREE-äge™ trademark of Arborjet, Inc.

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

Label Code: PARGNO016AA0109